

Josephson tunneling in high- T_c superconductors

M. B. Walker and J. Luetttmer-Strathmann

Department of Physics, University of Toronto, Toronto, Ontario, M5S 1A7, Canada

(received: December 19, 1995)

This article describes the Josephson tunneling from time-reversal symmetry-breaking states and compares it with that from time-reversal invariant states for both twinned and untwinned crystals and for both c -axis and basal-plane currents, in a model for orthorhombic YBCO. A macroscopic invariance group describing the superconducting state of a twinned crystal is introduced and shown to provide a useful framework for the discussion of the results for twinned crystals. In addition, a ring geometry, which allows s -wave and $d_{x^2-y^2}$ -wave superconductivity in a tetragonal superconductor to be distinguished on the basis of symmetry arguments only, is proposed and analyzed. Finally, an appendix gives details of the experimental Josephson tunneling evidence for a superconducting state of orthorhombic $ux^2 + vy^2$ symmetry in YBCO.

74.20.De, 74.50.+r, 74.72.Bk, 74.62.Bf

I. INTRODUCTION

Recently a number of Josephson tunneling experiments^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15} have been performed on the high- T_c material YBCO to investigate the symmetry of its order parameter. Conventional superconductors all have order parameters with s -wave symmetry. On the other hand, many theoretical models of high temperature superconductivity based on a description of the pairing of the electrons in approximately tetragonal copper-oxide layers (such as occur in YBCO) find superconducting order parameters of symmetry $d_{x^2-y^2}$ (see Refs. 16 and 17 for reviews). Thus, experiments have tended to focus on distinguishing between s and $d_{x^2-y^2}$ symmetries for the superconducting order parameter.

It is of interest to note that this focus on the YBCO order parameter as a linear combination of $d_{x^2-y^2}$ and s -wave order parameters can also be arrived at from a study of existing experimental results only, without appeal to current microscopic theories. If the transition to the superconducting state is continuous, then according to the Landau theory of second-order phase transitions, the superconducting order parameter must transform according to one of the eight irreducible representations of the orthorhombic point group of YBCO. It is shown in Appendix A, that the observation of nonzero Josephson tunneling currents along all three principal crystallographic directions in YBCO (e.g. as in Refs. 1, 2, 11) is sufficient to establish that the YBCO superconducting order parameter is a linear combination of $d_{x^2-y^2}$ - and s -wave components.

An important concept developed in this paper is that of the macroscopic symmetry (i.e. macroscopic invariance group) for the superconducting state of a twinned crystal. This macroscopic invariance group (which is different from the microscopic invariance group describing the superconducting symmetry of an untwinned single crystal) is the symmetry group determined by Josephson tunneling experiments on twinned crystals. The majority^{1,3,4,5,6,7,8,9,10} of Josephson tunneling experiments on twinned YBCO crystals have determined the macroscopic symmetry of the superconducting state to be the same as that of a tetragonal $d_{x^2-y^2}$ superconductor.

According to Ref. 18 (see also Ref. 19), two types of twin boundaries are possible within a Ginzburg-Landau model of YBCO in which the individual twins are in $d_{x^2-y^2} + s$ states: the superconducting state can have either even or odd reflection symmetry with respect to the twin boundary. Ref. 18 has shown that the experimental result of macroscopic $d_{x^2-y^2}$ superconductivity in twinned samples is equivalent to determining that the twin boundaries in YBCO have odd reflection symmetry.

This leaves the question of the relationship between microscopic and macroscopic symmetry, which we discuss briefly below. As it turns out, a microscopic understanding of the energetics of the twin boundaries is necessary to determine whether or not macroscopic $d_{x^2-y^2}$ symmetry implies a microscopic symmetry which is predominantly d -wave (see also Ref. 5). In this article, however, we prefer not to rely on microscopic arguments, and to consider all possibilities consistent with the requirements of symmetry.

In principle, (e.g. assuming that difficulties concerning trapped flux can be satisfactorily eliminated) a direct experimental determination of whether $d_{x^2-y^2}$ or s component of the order parameter is dominant can be made by performing a SQUID experiment on an untwinned single crystal of YBCO in the geometry suggested by Sigrist and Rice¹⁷. Experiments on untwinned crystals have been performed by Wollman *et al.*^{1,4,5} and by Brawner and Ott², and indicate that the microscopic symmetry of the order parameter is predominantly d -wave.

The tricrystal ring experiments of Tsuei *et al.*^{7,8,9,10,20} are of particular interest because of the precision with which problems associated with trapped flux can be eliminated, and because one of these experiments²⁰ has been carried out on $\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$, which is tetragonal and therefore not open to questions of interpretation related to the existence of twins. These experiments show convincingly rings containing a flux of one half an elementary flux quantum in their ground state; such rings must contain either one or three π Josephson junctions. Based on simplified models for the angular dependence (on the angles between the Josephson junction and the crystallographic axes) of the Josephson current, a plausible argument is made that these results imply $d_{x^2-y^2}$ superconductivity. However, it would not contradict the basic principles of physics if, for the experimental geometries used up to the present, s -wave superconductivity were to give rise to a ring whose ground state contained half an elementary flux quantum. We therefore propose and analyze, in a section of our article below, a ring geometry where the observation of one half a flux quantum in the ground state can exclude, on the basis of experimental measurements and symmetry arguments only, the possibility that the superconducting state is s -wave. The case of $d + is$ superconductivity (see below) is also discussed for this geometry.

The experiments of Ref. 15 do not appear to yield the $d_{x^2-y^2}$ macroscopic symmetry obtained by other Josephson experiments performed on twinned crystals. A possible explanation for this apparent inconsistency has been given in Ref. 21 in terms of the existence of vortices in the weak links. Further study of this experimental configuration has indeed found magnetic flux trapped in the grain boundary junctions²².

A remaining puzzle among the Josephson results on YBCO-Pb junctions concerns the c -axis critical currents. Because of the orthorhombicity of YBCO, there should be a nonzero c -axis Josephson current from an untwinned single crystal even if the superconducting order parameter of YBCO is predominantly $d_{x^2-y^2}$ -wave. In the measurements of Ref. 11, 12, 13, the c -axis current was found to be nonzero both for twinned and untwinned crystals of YBCO; the $I_c R_n$ product (where I_c is the critical current and R_n is the normal state resistance) is approximately an order of magnitude smaller for films than it is for single crystals¹², with the highest value being obtained for untwinned single crystals¹³. On the other hand, efforts¹⁴ to detect Josephson tunneling in cases where the Josephson junctions have a relatively high value of R_n have not been successful, perhaps due to thermal fluctuation effects. These varied results are not understood. According to Ref. 18, experimental measurements of basal-plane Josephson currents indicate that the twin boundaries have odd reflection symmetry; for such twin boundaries, the c -axis Josephson currents from twinned crystals will average to zero, assuming the two types of twins are present with equal weight. One possibility¹⁸ of accounting for a nonzero c -axis current is that the twins are not present with equal weight (perhaps as a result of strains); alternatives will be discussed below.

It is perhaps useful to introduce the idea of “robustness” of the consequences of the macroscopic symmetry. The macroscopic invariance group of a twinned crystal is determined by assuming that all types of twins are present with equal weight. However, it may happen that, due to strain for example, imbalances occur where one type of twin has greater weight than another. A robust property is one which is not changed by a small imbalance of this type. It can be shown that the observation of a π phase shift in a basal-plane corner SQUID experiment on YBCO is a robust property, whereas the expected zero value of the c -axis Josephson current from a twinned sample having macroscopic $d_{x^2-y^2}$ symmetry is not. Thus, as noted in the previous paragraph, the c -axis results of Refs. 11, 12, 13 could be explained in terms of an unequal weighting of twin types.

As noted by Li *et al.*²³ in a Ginzburg-Landau study of mixed s and $d_{x^2-y^2}$ superconductivity in an orthorhombic superconductor such as YBCO, the phase transition to the superconducting state will be, assuming that this transition is continuous, to a time-reversal invariant state called the $d + s$ state. They also pointed out²³ that a second transition to a different superconducting state called the $d + e^{i\phi}s$ state is possible at a lower temperature. This $d + e^{i\phi}s$ state breaks time reversal symmetry. At present there is only one report²⁴ of a second phase transition in YBCO. It is however possible, particularly in twinned samples, that such a second transition could be smeared out and not easily visible. The $d + e^{i\phi}s$ state is not stable in some microscopic theories²⁵; a recent weak-coupling calculation²⁵, for example, finds that the $d + s$ is always stable relative to the $d + e^{i\phi}s$ state. Since there is at present no consensus concerning an appropriate microscopic theory of high temperature superconductivity, we do not exclude the $d + e^{i\phi}s$ state from our considerations but explore its experimental consequences.

This article begins by discussing the time-reversal symmetry-breaking $d + e^{i\phi}s$ state, the possible types of twin boundaries consistent with this state, and the macroscopic symmetries of twinned crystals having twins in the $d + e^{i\phi}s$ state. Some results relevant to $d + s$ states, which are a special case ($\phi = 0$) are included for completeness.

Broken time-reversal symmetry in YBCO has been searched for in optical experiments^{26,27}, μSR experiments²⁸, and in Josephson tunneling experiments³ and has not been found. Nevertheless, because of the importance of the Josephson effect as a method of determining the order parameter symmetry, it is essential to have a clear understanding of the experimental manifestations of broken time-reversal symmetry for experiments performed on both untwinned and twinned crystals. In related work, some effects of broken time-reversal symmetry at crystal grain boundaries²⁹ and at twin boundaries³⁰ have recently been considered by other authors.

The Josephson current through a junction can be written as

$$I = I_c \sin(\phi - \phi_c), \quad (1.1)$$

where $I_c > 0$ is the Josephson critical current, and ϕ is the gauge invariant difference between the phases of the superconductors on the two sides of the junction. For normal junctions (e.g. junctions between conventional s -wave superconductors) the characteristic phase of the junction can always be taken to be $\phi_c = 0$. Experiments^{1,2,3,4,5,6,7,8,9,10,20} on high temperature superconductors on the other hand, give evidence for circuits containing an odd number of π junctions, i.e. junctions having a characteristic phase of $\phi_c = \pi$. The calculations performed below show, for both twinned and untwinned crystals, that a Josephson junction involving a superconductor in a state which breaks time-reversal symmetry can have a characteristic phase ϕ_c which is neither zero nor π . A similar conclusion concerning ϕ_c was previously reached in Ref. 8.

In sections V and VI, the Josephson currents for twinned samples are calculated by averaging over currents from the individual twins; the result of this averaging is shown to be consistent with the expectations based on the appropriate macroscopic invariance group for the twinned sample.

In this article, it is assumed that the orthorhombic basal-plane unit cells of neighboring twins are at right angles to each other, cf. Fig. 1, whereas in fact they are rotated towards each other by a small angle. This misorientation was considered carefully in Ref. 18 and can be shown not to alter the results for Josephson tunneling discussed here.

II. TWIN BOUNDARIES BETWEEN STATES BREAKING TIME-REVERSAL SYMMETRY

The Landau free energy density describing coupled s - and $d_{x^2-y^2}$ -wave superconductivity in an orthorhombic superconductor such as YBCO is²³

$$\begin{aligned} f_L = & \alpha_s |\psi_s|^2 + \alpha_d |\psi_d|^2 \\ & + \frac{1}{2} \beta_s |\psi_s|^4 + \frac{1}{2} \beta_d |\psi_d|^4 + \beta_4 |\psi_s|^2 |\psi_d|^2 \\ & + (-1)^\epsilon \alpha (\psi_s \psi_d^* + \psi_s^* \psi_d) + \beta_0 (\psi_s^2 \psi_d^{*2} + \psi_s^{*2} \psi_d^2) \\ & + (-1)^\epsilon [\beta_1 |\psi_d|^2 + \beta_2 |\psi_s|^2] (\psi_s \psi_d^* + \psi_s^* \psi_d), \end{aligned} \quad (2.1)$$

where ψ_s and ψ_d denote the s - and $d_{x^2-y^2}$ -wave components of the order parameter, respectively, where $\alpha_s = \alpha'_s (T - T_{s0})$, $\alpha_d = \alpha'_d (T - T_{d0})$, $\alpha'_s, \alpha'_d > 0$, T is the temperature, and where $\epsilon \in \{1, 2\}$ indicates which of the two types of neighboring twins is described, cf. Fig. 1. The appearance of the $(-1)^\epsilon$ factors in Eq. (2.1), which is a generalization of the free energy presented in Ref. 23, is related to the assumption that the twin boundary between two twins is a plane of reflection symmetry of the underlying crystal lattice¹⁸. With the definition of the coordinates in Fig. 1, the $d_{x^2-y^2}$ component of the order parameter, ψ_d , changes sign under reflection in the twinning plane. To ensure that the free energy of the reflected state is the same as that of the original state factors of $(-1)^\epsilon$ are included in Eq. (2.1).

For $\alpha = \beta_1 = \beta_2 = 0$, with $\beta_s > 0$, $\beta_d > 0$, and $\beta_s \beta_d - (\beta_4 - 2|\beta_0|)^2 > 0$ for thermodynamic stability, the free energy (2.1) describes a tetragonal system undergoing a continuous phase transition at $T = \max(T_{d0}, T_{s0})$ to either a d -wave ($\psi_d \neq 0$, $\psi_s = 0$) or an s -wave ($\psi_s \neq 0$, $\psi_d = 0$) superconducting state. A second transition to a mixed ($\psi_d \neq 0$, $\psi_s \neq 0$) state occurs when $\alpha_s \beta_d = \alpha_d (\beta_4 - 2|\beta_0|)$ or $\alpha_d \beta_s = \alpha_s (\beta_4 - 2|\beta_0|)$, respectively. We define the relative phase ϕ_{sd} between ψ_s and ψ_d through

$$\psi_d = |\psi_d| e^{i\phi_d}, \quad \psi_s = |\psi_s| e^{i(\phi_d + \phi_{sd})}. \quad (2.2)$$

In the tetragonal case, ϕ_{sd} is determined by the sign of β_0 : $\phi_{sd} \in \{0, \pi\}$ for $\beta_0 < 0$, corresponding to a time-reversal invariant $d + s$ state, $\phi_{sd} \in \{\pi/2, 3\pi/2\}$ for $\beta_0 > 0$, corresponding to a time-reversal symmetry-breaking $d + is$ state.

The terms with coefficients α , β_1 , and β_2 describe the coupling of the s - and d -wave components of the order parameter which exists in an orthorhombic crystal. A continuous phase transition from the normal to a superconducting state occurs in the orthorhombic system when the smallest eigenvalue of the quadratic form in Eq. (2.1) goes through zero. For $\alpha \ll 1$, i.e. weak orthorhombicity, and $\alpha_d < \alpha_s$ the transition is at temperature $T_d \approx T_{d0} + \alpha^2 / \alpha_s \alpha'_d$ to a state whose d -wave component ψ_d grows with decreasing temperature approximately like $|\psi_d|^2 = \alpha'_d (T_d - T) / \tilde{\beta}_d$, where $\tilde{\beta}_d \approx \beta_d$. The s -wave component ψ_s can be found as an expansion in powers of ψ_d , and to lowest order is

$$\psi_s = (-1)^{\epsilon+1} (\alpha / \alpha_s) \psi_d. \quad (2.3)$$

For this solution, ψ_d and ψ_s have the same phase (modulo π) and the superconducting state is called a $d + s$ state. For $|\alpha / \alpha_s| \ll 1$, $|\psi_s| \ll |\psi_d|$, and the superconducting state is said to be predominantly d -wave. For $\alpha_s < \alpha_d$, on the other hand, a transition occurs at $T_s \approx T_{s0} + \alpha^2 / \alpha_d \alpha'_s$ to a predominantly s -wave superconducting state.

To determine the relative phase ϕ_{sd} between ψ_s and ψ_d , cf. Eq. (2.2), we set $\alpha > 0$ and, for simplicity, extend the assumption of weak orthorhombicity by neglecting β_1 and β_2 in comparison with β_0 . By minimizing the free energy (2.1) we find, for $\beta_0 > 0$, the following conditions on the relative phase ϕ_{sd} :

$$\sin \phi_{sd} = 0, \quad (2.4)$$

or

$$\cos \phi_{sd} = (-1)^{\epsilon+1} \frac{\alpha}{4\beta_0 |\psi_s| |\psi_d|}. \quad (2.5)$$

Condition (2.4) yields a relative phase of $\phi_{sd} = 0$ for $\epsilon = 1$, $\phi_{sd} = \pi$ for $\epsilon = 2$, and a time-reversal invariant $d + s$ superconducting state. This state is realized near the transition from the normal state (see also Eq. (2.3)) and, as the temperature is lowered, remains stable until $|\psi_d|$ and $|\psi_s|$ have grown sufficiently to reduce the fraction on the right-hand side of Eq. (2.5) to unity. At the temperature $T = T_\phi$ with $|\psi_s(T_\phi)| |\psi_d(T_\phi)| = \alpha/4\beta_0$ a second transition to a $d + e^{i\phi}s$ state takes place. For the $\epsilon = 1$ twin, for example, $|\phi_{sd}|$ increases continuously from zero for temperatures below T_ϕ . The values $\phi_{sd} = \pm |\phi_{sd}|$ correspond to two energetically equivalent states of twin $\epsilon = 1$ that are related to each other by the time-reversal operation $\mathcal{T}$. The $d + e^{i\phi}s$ state thus breaks time-reversal symmetry.

In the case that $\beta_0 < 0$, on the other hand, the minimum of the free energy (2.1) is always attained for $\phi_{sd} = 0$ for $\epsilon = 1$ and $\phi_{sd} = \pi$ for $\epsilon = 2$, so that only one phase transition occurs and the superconducting state is always the time-reversal invariant state $d + s$.

Our considerations so far have shown that the relative phase ϕ_{sd} between the components of the order parameter in a given twin is determined by the bulk free energy density (2.1). Contributions to the free energy due to the interaction between the order parameters on opposite sides of a twin boundary have to be considered to describe the relative phases of the order parameter components across a twin boundary. For a predominantly d -wave superconductor the interface free energy per unit area associated with interactions between twin 1 and twin 2, cf. Fig. 1, is to lowest order given by¹⁸

$$f_{12}^d = B(\psi_{d1}^* \psi_{d2} + \psi_{d1} \psi_{d2}^*), \quad (2.6)$$

where ψ_{d1} and ψ_{d2} denote the d -wave components of the order parameter in twin 1 and 2, respectively. For time-reversal invariant $d + s$ states, the free energy is minimized by $\psi_{d1} = \psi_{d2}$ ($\psi_{d1} = -\psi_{d2}$) for $B < 0$ ($B > 0$), which implies that the overall superconducting states have the form

$$\Psi_u = (\psi_{d1}, \psi_{s1}, \psi_{d2}, \psi_{s2}) = (\psi_d, \psi_s, \psi_d, -\psi_s), \quad B < 0, \quad (2.7)$$

and

$$\Psi_g = (\psi_{d1}, \psi_{s1}, \psi_{d2}, \psi_{s2}) = (\psi_d, \psi_s, -\psi_d, \psi_s), \quad B > 0. \quad (2.8)$$

The states Ψ_g and Ψ_u satisfy $\sigma \Psi_g = \Psi_g$ and $\sigma \Psi_u = -\Psi_u$ and are thus even and odd with respect to a reflection σ in the twinning plane, cf. Fig. 2. An investigation of Josephson tunneling currents for time-reversal invariant states has shown¹⁸ that the experimental results for YBCO are consistent with odd symmetry under σ , so that the d -wave component is the same on either side of the twin boundary.

The assumption that $\psi_{d1} = \psi_{d2}$ remains correct for the time-reversal symmetry-breaking $d + e^{i\phi}s$ states yields two possible states in twin 2 for a given state in twin 1:

$$\Psi_1 = e^{i\phi_d}(|\psi_d|, e^{i\phi_{sd}}|\psi_s|), \quad (2.9)$$

$$\Psi_2 = e^{i\phi_d}(|\psi_d|, -e^{i\phi_{sd}}|\psi_s|), \quad (2.10)$$

$$\Psi_{\bar{2}} = e^{i\phi_d}(|\psi_d|, -e^{-i\phi_{sd}}|\psi_s|). \quad (2.11)$$

Since $\Psi_2 = -\sigma \Psi_1$, and since the state $\Psi_{\bar{2}}$ is related to the time reversed state of twin 1, defined as $\Psi_{\bar{1}} = e^{i\phi_d}(|\psi_d|, e^{-i\phi_{sd}}|\psi_s|)$, by $\Psi_{\bar{2}} = -\sigma \Psi_{\bar{1}}$, it is clear that all states have the same bulk free energy density. Which of these two states of twin 2 prefers to be in contact with the given state of twin 1 is determined by the surface energy of interaction of twin 1 with twin 2. Note that if the pair (Ψ_1, Ψ_2) is favored, the overall state is odd with respect to a reflection σ in the twinning plane; this type of twin boundary will be called an odd-symmetry twin boundary. Otherwise, if the pair $(\Psi_1, \Psi_{\bar{2}})$ is favored, the overall superconducting state is invariant, to within a phase factor, with respect to a reflection σ followed by time reversal $\mathcal{T}$; this second type of twin boundary will be called a $\mathcal{T}\sigma$ twin boundary. For a given material, only one type of twin boundary will be present; this will be the one which contributes the least to the free energy.

III. MACROSCOPIC SYMMETRY

Since a twinned sample will have some regions in which the twinning planes are (110) planes, and other regions where the twinning planes are ($\bar{1}10$) planes, it is important to examine how the superconducting order is transferred from one region to another. Consider the case where the twinning planes are odd-symmetry reflection planes. Within a given twin, say twin 1, the superconductivity is described by its order parameter Ψ_1 , which is a mixture of $d_{x^2-y^2}$ - and s -wave components (cf. Eq. (2.9)). This order parameter is invariant with respect to the orthorhombic point group operations (rotations of π about the orthorhombic principal axes and reflections in planes normal to the principal axes) and hence, from the point of view of its point group symmetry, can be represented by a rectangle, as in Fig. 3. (This same rectangle can also be used to indicate the relative orientation of the a and b axes in the different twins.) At an odd-symmetry twin boundary, not only does the rectangle representing the order parameter change its orientation on reflection across a twin boundary, but it changes sign. Thus, in Fig. 3, the alternating plus and minus signs in the rectangles represent the different signs of the order parameters in the different twins. Also, the interface between a region of (110) twin boundaries and a region of ($\bar{1}10$) twin boundaries, as determined by experiment³¹, is shown in Fig. 3. It is of interest that this interface is itself largely a twin boundary, with small breaks in this boundary. Thus, the superconducting order parameter should be transmitted from a region of (110) twin boundaries to a region of ($\bar{1}10$) twin boundaries in exactly the same way as it is transmitted from one twin to another within a region of, say, (110) twin boundaries. In the case of $\mathcal{T}\sigma$ twin boundaries, the superconducting order is passed from twin to twin and between regions of differently oriented twin boundaries in a similar way.

A. Macroscopic invariance group for $d + e^{i\phi}s$ states

Figure 3 also gives a simple picture of the macroscopic symmetry of a highly twinned superconductor with odd-symmetry twin boundaries. Note that Fig. 3(b) is obtained from Fig. 3(a) by a rotation of $\frac{1}{2}\pi$ about the c -axis. However, Fig. 3(b) is also obtained from Fig. 3(a) by changing the signs of all of the order parameters, i.e. by multiplying the state of the entire system by minus one. Thus, a rotation of this twinned crystal by $\frac{1}{2}\pi$ about the c -axis is equivalent to changing the sign of the overall order parameter. In other words, the superconducting state of the twinned crystal is invariant under the operation $e^{i\pi}C_4$. Similarly, the superconducting state of the twinned crystal can be shown to be invariant with respect to all of the operations of the direct product group $D_4^{(1)}(D_2) \times I$, where I is the group containing the identity and the inversion, and $D_4^{(1)}(D_2)$ is the combined group $D_4^{(1)}(D_2) = (E, C_2, 2U_2, 2C_4e^{i\pi}, 2U_2'e^{i\pi})$ in the notation of Ref. 32. The direct product group $D_4^{(1)}(D_2) \times I$ will be said to describe the macroscopic symmetry of superconductivity in the twinned crystal. Note that this macroscopic symmetry group lacks time reversal; the time-reversal operation creates a different superconducting state with the same free energy.

The macroscopic invariance group, $D_4^{(1)}(D_2) \times I$, of a twinned crystal with twins in $d + e^{i\phi}s$ states and with odd reflection-symmetry twin boundaries is thus the same, except for the lack of the time-reversal operation, as the invariance group of the $d_{x^2-y^2}$ superconducting state of an (untwinned) tetragonal single crystal³². If the characteristics of the Josephson effect are determined by the macroscopic invariance group of the twinned crystal, then we expect the c -axis Josephson current to be zero, and an offset flux (cf. Eq. (6.2) below) of $\frac{1}{2}\Phi_0$ in a corner SQUID experiment. These characteristics of the Josephson effect for the type of twinned-crystal superconductivity described in this paragraph are confirmed by detailed twin averaging calculations in Sections V and VI.

The discussion of the case where the superconducting twin boundaries are of type $\mathcal{T}\sigma$ can be carried out in a similar way. In this case, the basic states of the two types of twins are Ψ_1 and Ψ_2 , as given by Eqs. (2.9) and (2.11). Detailed consideration of the effects of the elements of D_4 shows that the macroscopic invariance group of this state of the twinned crystal is $D_4^{(3)}(D_2) \times I$, where $D_4^{(3)}(D_2) = (E, C_2, 2U_2, 2\mathcal{T}e^{i\pi}C_4, 2\mathcal{T}e^{i\pi}U_2')$. Thus, the macroscopic symmetry here is the same as that of the $d + is$ superconducting state of an (untwinned) tetragonal crystal. Again, the time-reversal operation $\mathcal{T}$ by itself does not leave the state invariant, but rather creates a different state with the same free energy. The macroscopic symmetry here allows a nonzero c -axis Josephson current and an offset flux differing from $\frac{1}{2}\Phi_0$ in a corner SQUID experiment; detailed calculations performed below indeed confirm these properties for a twinned crystal with $\mathcal{T}\sigma$ twin boundaries.

B. Macroscopic invariance group for $d + s$ states

A previous paper¹⁸ which presented detailed calculations of the Josephson currents from an orthorhombically twinned crystal, with twins in the time-reversal invariant $d + s$ state, to a conventional superconductor, did not

exploit the idea of the macroscopic symmetry of the twinned crystal. Two types of twin boundaries are possible, giving rise to superconducting states that are either odd, cf. Eq. (2.7), or even, cf. Eq. (2.8), with respect to a reflection in the twinning plane. The case of odd reflection-symmetry twin boundaries is the same as the case of odd reflection-symmetry twin boundaries studied above, except that $\phi_{sd} = 0$ since the twins are in time-reversal invariant states. Hence, the time-reversal operation $\mathcal{T}$ is added to the invariance group, which becomes $D_4^{(1)}(D_2) \times I \times R$, where R contains the identity and time-reversal; this is also the invariance group for $d_{x^2-y^2}$ superconductivity in a tetragonal superconductor. For the case of even reflection-symmetry twin boundaries, which can be represented schematically by assigning plus signs to all of the rectangles in Fig. 3, the macroscopic invariance group is $D_4 \times I \times R$, which is also the invariance group for s -wave superconductivity in a tetragonal superconductor. Thus, in orthorhombically twinned crystals, with the twins in $d + s$ superconducting states, odd reflection symmetry twin boundaries give a macroscopic symmetry identical to that of a tetragonal $d_{x^2-y^2}$ -wave superconductor, whereas even reflection-symmetry twin boundaries give a macroscopic symmetry identical to that of a tetragonal s -wave superconductor. Clearly, as emphasized in Ref. 18, the qualitative characteristics of the Josephson effect in twinned crystals, which are determined by the macroscopic symmetry of the superconducting state, are a reflection of the symmetry of the twin boundaries, and have nothing to do with whether the basic $d + s$ state of the twins is predominantly $d_{x^2-y^2}$ -wave ($|\psi_d| \gg |\psi_s|$) or predominantly s -wave ($|\psi_s| \gg |\psi_d|$).

C. Relation between twin-boundary symmetry and microscopic symmetry

Given that Josephson experiments on twinned crystals of YBCO determine directly the twin boundary symmetry, it is of interest to ask to what extent this result can be used to infer the dominant microscopic order-parameter symmetry.

As noted above, if a phenomenological Ginzburg-Landau type of model is adopted for the twin boundaries (e.g. as in Eq. (2.6) above or as in Ref. 19) they can have either even or odd reflection symmetry. Which case occurs depends on the signs and magnitudes of certain parameters in the twin-boundary free energy. The fact that the microscopic order parameter may be either predominantly $d_{x^2-y^2}$ -wave or predominantly s -wave places no restrictions on these parameters at the phenomenological level. Thus, within the framework of Ginzburg-Landau type models, the identification of the twin-boundary symmetry does not lead to information about the dominant order parameter symmetry.

Now consider the odd-symmetry and even-symmetry twin boundaries as shown in Fig. 2. For the odd-symmetry twin boundary, the s -wave component of the order-parameter changes sign, whereas the d -wave component is continuous. On the other hand, for an even-symmetry twin boundary, the d -wave component of the order parameter changes sign, while the s -wave component is continuous. If it is assumed that either the $d_{x^2-y^2}$ - or s -wave component of the order parameter is completely dominant, and that the lowest free energy state is achieved by having an order parameter that is continuous and varies as little as possible at the twin boundary, then predominantly $d_{x^2-y^2}$ -wave (s -wave) superconductivity would lead to odd-symmetry (even symmetry) twin boundaries. The argument that the dominant order parameter should be as continuous as possible seems reasonable, but there are at least hypothetical situations (see e.g. Ref. 33) for which it is not correct.

In YBCO, both infrared³⁴ and Josephson tunneling³⁵ measurements show a relatively strong anisotropy in the ab -plane penetration depth. Furthermore, a recent microscopic analysis³⁶ of the penetration depth data indicates that the CuO chains play a significant role in the superconductivity. This suggests that neither the $d_{x^2-y^2}$ nor the s component of the order parameter is completely dominant, but that both have significant amplitudes. For an odd-symmetry twin boundary, the assumption that the order parameters vary continuously across the twin boundary implies that the $d_{x^2-y^2}$ -wave component of the order parameter remains approximately uniform across the twin boundary. The s -wave component, however, will have its bulk value ψ_s at distances greater than $-\xi_s$ from the twin boundary (ξ_s is the s -wave coherence length), will go continuously to zero and change sign at the twin boundary, and will tend to the value $-\psi_s$ at distances greater than $+\psi_s$ from the twin boundaries.

If the condensation energy per unit volume gained by having a nonzero s -wave order parameter is ϵ_s , then the free energy per unit area of an odd reflection-symmetry twin boundary will, in this model, be approximately

$$f_{\text{odd}} = 2\epsilon_s\xi_s + \delta f_{\text{odd}}. \quad (3.1)$$

Here a quantity δf_{odd} is assumed to include all corrections necessary to arrive at the correct answer; these corrections may be due to interactions between the s and d -wave components of the order parameter, effects of crystal structure and doping level at the twin boundary giving condensation energies not equal to the bulk value, the effect of disorder at the twin boundary which might reduce the amplitude of the d -wave component of the order parameter, etc.

Similarly, in the case of an even reflection-symmetry twin boundary, the twin-boundary free energy per unit area for our simplified model can be written

$$f_{\text{even}} = 2\epsilon_d \xi_d + \delta f_{\text{even}}. \quad (3.2)$$

Within this basic model of the energetics of the twin boundary, a knowledge of the relative condensation energies, and of the relative coherence lengths of the s - and $d_{x^2-y^2}$ -wave components of the order parameter is required to determine whether the odd-symmetry or even-symmetry twin boundary is stable for a given relative weight of s and $d_{x^2-y^2}$ to the order parameter. We are not aware of the existence of good quantitative estimates of these quantities.

In summary, our view is that, while it may be possible to make appropriate assumptions which would suggest that the observation of odd-symmetry twin boundaries implies a dominant $d_{x^2-y^2}$ component of the order parameter, the argument is not conclusive. There is thus an important role for experiments on untwinned single crystals of YBCO^{1,2,4,5}, or on tetragonal crystals such as $\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$ ²⁰, since in these cases the complications due to twinning are eliminated.

Finally, it should be noted that related issues have also been discussed in Ref. 5, where a somewhat different point of view is taken.

IV. JOSEPHSON CURRENTS

This section establishes a relationship between the Josephson current across a junction and the phase derivative of the free energy by extending the approach of Andreev¹⁹ (see Ref. 17 for a review) to the case of a multicomponent order parameter. In order to calculate currents in the superconductor, gradient terms compatible with the orthorhombic symmetry of the system are added to the free energy density (2.1). To lowest order, these are given by

$$F_{\text{grad}} = \int dV \frac{1}{2} \sum_{\alpha} \sum_{\lambda, \mu} \frac{1}{m_{\lambda, \mu}^{\alpha}} (p_{\alpha} \psi_{\lambda})^* (p_{\alpha} \psi_{\mu}), \quad (4.1)$$

where $\alpha \in \{x, y, z\}$ and $\lambda, \mu \in \{d, s\}$. The coefficients of the gradient terms are real, satisfy $m_{\lambda, \mu}^{\alpha} = m_{\mu, \lambda}^{\alpha}$, and the components p_{α} of the “momentum” operator are given by

$$p_{\alpha} = -i\hbar \nabla + \frac{|e^*|}{c} A_{\alpha}, \quad (4.2)$$

where A_{α} is a component of the vector potential, c is the speed of light, and $|e^*|$ is the absolute value of the charge of a Cooper pair. The components j_{α} of the current are obtained from the variation of F_{grad} with respect to A_{α} , and are given by

$$j_{\alpha} = \sum_{\lambda, \mu} \frac{|e^*|}{2m_{\lambda, \mu}^{\alpha}} \psi_{\lambda}^* p_{\alpha} \psi_{\mu} + \text{c.c.} \quad (4.3)$$

To derive a Josephson current-phase relation for our unconventional superconductor, we generalize definition (2.2) by writing $\psi_{\mu} = |\psi_{\mu}| e^{i(\phi_d + \phi_{\mu d})}$, where $\phi_{dd} = 0$, and ϕ_{sd} is the relative phase defined in Eq. (2.2). The phase ϕ_d can vary with position but, as pointed out in the previous section, ϕ_{sd} is determined by the bulk free energy. Under the assumption that $|\psi_{\mu}|$ is independent of position, we find:

$$j_{\alpha} = c_{\alpha} \left[\frac{\partial \phi_d}{\partial x_{\alpha}} + \frac{2\pi}{\Phi_0} A_{\alpha} \right], \quad \Phi_0 = \frac{hc}{|e^*|}, \quad (4.4)$$

where

$$c_{\alpha} = \sum_{\lambda, \mu} \frac{\hbar |e^*|}{m_{\lambda, \mu}^{\alpha}} |\psi_{\lambda}| |\psi_{\mu}| \cos(\phi_{\mu d} - \phi_{\lambda d}). \quad (4.5)$$

The Josephson current across an interface between YBCO and a conventional superconductor like lead depends on the interface free energy associated with interactions between the s - and d -wave components $\psi_{\mu}, \mu \in \{s, d\}$ of the order parameter on the YBCO side and the conventional (s -wave) order parameter ψ_{PB} on the lead side. This interface free energy per unit area can be written as:

$$f_I = \sum_{\mu} G_{\mu}(\mathbf{n}) (\psi_{\mu}^* \psi_{\text{Pb}} + \text{c.c.}), \quad (4.6)$$

where $G_{\mu}(\mathbf{n})$ is real and, in general, depends on the direction of the surface normal $\mathbf{n}$, relative to the YBCO crystallographic axes. There are also contributions to the interface free energy that depend on ψ_{μ} and ψ_{Pb} separately. Since these terms do not contribute to the tunneling we neglect them here.

The minimization of the total interface free energy with respect to variations in ψ_{λ}^* yields

$$\sum_{\alpha, \mu} n_{\alpha} \frac{i\hbar}{2m_{\lambda, \mu}^{\alpha}} (p_{\alpha} \psi_{\mu}) = -G_{\lambda}(\mathbf{n}) \psi_{\text{Pb}}. \quad (4.7)$$

Insertion of Eq. (4.7) into (4.3) gives, for the Josephson current $\mathbf{j}$ across an interface between YBCO and lead with normal vector $\mathbf{n}$,

$$\mathbf{n} \cdot \mathbf{j} = \frac{-i|e^*|}{\hbar} \sum_{\lambda} G_{\lambda}(\mathbf{n}) \psi_{\lambda}^* \psi_{\text{Pb}} + \text{c.c.} \quad (4.8)$$

As in the case of conventional superconductivity, the current $\mathbf{j}$ is related to the derivative of the interface free energy f_I with respect to the phase difference between the order parameters on both sides of the junction. Substituting $\psi_{\text{Pb}} = |\psi_{\text{Pb}}| e^{i\phi_{\text{Pb}}}$, and ψ_d and ψ_s from Eq. (2.2) into Eqs. (4.6) and (4.8), and setting $\phi = \phi_d - \phi_{\text{Pb}}$ yields

$$\frac{\partial f_I(\phi)}{\partial \phi} = \frac{\Phi_0}{2\pi c} \mathbf{n} \cdot \mathbf{j}. \quad (4.9)$$

V. c-AXIS TUNNELING

Consider as a first example Josephson tunneling along the c -axis between YBCO and lead. The free energy per unit area f_I describing the interaction between the superconducting order parameters at the interface, which is normal to the c -axis, can be written as:

$$f_I = - [(-1)^{\epsilon+1} D \psi_d^* + S \psi_s^*] \psi_{\text{Pb}} + \text{c.c.}, \quad (5.1)$$

where the constants $(-1)^{\epsilon+1} D$ and S are appropriate realizations of the coefficients $G_{\mu}(\mathbf{n})$ in Eq. (4.6). The coefficient D is nonzero because of the orthorhombicity of YBCO.

We start with the interface energy $f_{I,1}$ for twin 1 for the case that YBCO is in the time-reversal symmetry-breaking $d + e^{i\phi} s$ state. Let twin 1 be in the superconducting state Ψ_1 defined in Eq. (2.9) and let $\psi_{\text{Pb}} = |\psi_{\text{Pb}}| e^{i\phi_{\text{Pb}}}$. With the definitions

$$d = 2D|\psi_d| |\psi_{\text{Pb}}|, \quad s = 2S|\psi_s| |\psi_{\text{Pb}}|, \quad \phi = \phi_d - \phi_{\text{Pb}}, \quad (5.2)$$

we find

$$f_{I,1} = -r_1 \cos(\phi - \phi_1), \quad (5.3)$$

with

$$r_1 = \sqrt{(d + s \cos \phi_{sd})^2 + (s \sin \phi_{sd})^2} \\ \sin \phi_1 = -s \sin \phi_{sd} / r_1, \quad \cos \phi_1 = (d + s \cos \phi_{sd}) / r_1. \quad (5.4)$$

Note that for the time-reversal symmetry-breaking state (i.e. $\phi_{sd} \neq 0$), the characteristic phase of the junction, ϕ_1 , will be neither 0 nor π .

If twin 2 is in the state $\Psi_2 = -\sigma \Psi_1$, cf. Eq. (2.10), so that the twin boundary is of the odd symmetry type, then the free energy $f_{I,2}$ follows from Eq. (5.3) for $f_{I,1}$ by replacing $\phi_{sd} \rightarrow \phi_{sd} + \pi$ and $d \rightarrow -d$ with the result:

$$f_{I,2} = -r_1 \cos(\phi - \phi_1 - \pi) = -f_{I,1}. \quad (5.5)$$

Hence, the average of the interface energy over the two types of twins $(f_{I,1} + f_{I,2})/2$ vanishes which implies, cf. Eq. (4.9), that the Josephson currents also average to zero. This result of the Josephson current averaging to zero for

odd symmetry twin boundaries is independent of the value of ϕ_{sd} and is thus the same for the time-reversal invariant and the time-reversal symmetry-breaking states. Hence, the result obtained above from the ideas of the macroscopic symmetry of a twinned crystal is confirmed: the element $e^{i\pi}C_4$ of the macroscopic invariance group will not allow c -axis Josephson currents.

Now suppose that twin 2 is in the state $\Psi_{\bar{2}} = -\sigma\Psi_{\bar{1}}$, cf. Eq. (2.11), so that the twin boundaries are of the $\mathcal{T}\sigma$ type. The free energy $f_{I,\bar{2}}$ is then obtained from Eq. (5.3) with the substitutions $\phi_{sd} \rightarrow -\phi_{sd} + \pi$ and $d \rightarrow -d$, which yields

$$f_{I,\bar{2}} = -r_1 \cos(\phi + \phi_1 - \pi). \quad (5.6)$$

The average of the interface free energies over the twins is then given by

$$\bar{f}_I = \frac{1}{2}(f_{I,1} + f_{I,\bar{2}}) = -r_1 \sin \phi_1 \sin \phi. \quad (5.7)$$

Together with Eq. (4.9) this yields for the current,

$$\mathbf{n} \cdot \mathbf{j} = -\frac{2\pi cr_1}{\Phi_0} \sin \phi_1 \cos \phi. \quad (5.8)$$

Hence, there is a partial cancellation of the Josephson currents upon averaging over twins such that the critical current j_c is reduced from the value for the untwinned crystal by a factor $|\sin \phi_1|$. This result of partial but not total cancellation of the Josephson currents in twinned crystals resembles to some extent what has been observed in experiments, namely, that the highest values of the $I_c R_n$ product have been observed in untwinned crystals¹³, and that smaller values (which are however nonzero) are observed in twinned crystals^{11,12,13} (see however Ref. 14). Note finally, that the macroscopic invariance group found above for the case of $\mathcal{T}\sigma$ twin boundaries allows a nonzero Josephson current, in agreement with what has been found here.

VI. BASAL PLANE JOSEPHSON CURRENTS

The experimental study of the basal plane Josephson currents is often carried out using the corner SQUID geometry^{1,2,3,17} of Fig. 4. The total current I_{total} , which is the sum of the currents through junctions A and B is

$$I_{\text{total}} = I_A \sin(\phi_A - \phi_{cA}) + I_B \sin(\phi_B - \phi_{cB}). \quad (6.1)$$

The difference in the gauge invariant phase differences for the two junctions is $\phi_A - \phi_B = 2\pi\Phi/\Phi_0$, where Φ is the flux through the SQUID and Φ_0 is the elementary flux quantum. The average phase $(\phi_A + \phi_B)/2$ is a free variable which adjusts itself so that the total current I_{total} equals the current fed into the SQUID. The maximum value of I_{total} with respect to variations of $(\phi_A + \phi_B)/2$ is

$$I_{\text{max}} = (I_A + I_B) \times \left\{ \epsilon^2 + (1 - \epsilon^2) \cos^2 \left[\frac{\pi}{\Phi_0} (\Phi - \Phi_{\text{offset}}) \right] \right\}^{1/2}, \quad (6.2)$$

where

$$\Phi_{\text{offset}} = \frac{\phi_{cA} - \phi_{cB}}{2\pi} \Phi_0, \quad (6.3)$$

and

$$\epsilon = (I_A - I_B)/(I_A + I_B). \quad (6.4)$$

Note that I_{max} is periodic in the flux Φ with period Φ_0 . The quantity Φ_{offset} is the flux at which I_{max} has its largest value. For two normal junctions $\Phi_{\text{offset}} = 0$, whereas for a SQUID with one normal and one π junction, Φ_{offset} has the value $\Phi_0/2$ ^{1,4,17}. It is now shown that Φ_{offset} can have other values when the superconducting state is not time-reversal invariant.

Consider a plane Josephson junction between twin 1 of YBCO and an s -wave superconductor such as lead. The junction is normal to the YBCO basal plane and its free energy $f_{I,1}$ per unit area is

$$f_{I,1}(\theta) = -[D(\theta)\psi_{d1}^* + S(\theta)\psi_{s1}^*]\psi_{Pb} + \text{c.c.}, \quad (6.5)$$

where θ is the angle between the normal to the interface $\mathbf{n}$ and the basis vector $\mathbf{a}_1$. By reflecting the state (ψ_{d1}, ψ_{s1}) of twin 1 with interface angle θ in the twin boundary, a state $(\psi_{d2}, \psi_{s2}) = (-\psi_{d1}, \psi_{s1})$ for twin 2 is obtained which has interface angle $-\theta + \pi/2$ and the same free energy. Hence the interface free energy for twin 2 is given by

$$f_{I,(2)}(\theta) = - \left[-D(-\theta + \frac{\pi}{2})\psi_{d2}^* + S(-\theta + \frac{\pi}{2})\psi_{s2}^* \right] \psi_{Pb} + \text{c.c.} \quad (6.6)$$

Orthorhombicity implies that

$$\begin{aligned} D(\theta + \pi) &= D(-\theta) = D(\theta), \\ S(\theta + \pi) &= S(-\theta) = S(\theta). \end{aligned} \quad (6.7)$$

With the aid of definition (2.9) for the superconducting state, the interface free energy for twin 1 can be written as

$$f_{I,1}(\theta) = -r(\theta) \cos(\phi - \phi_c(\theta)), \quad (6.8)$$

where $\phi = \phi_d - \phi_{Pb}$, and where

$$\begin{aligned} r(\theta) &= \sqrt{\tilde{d}^2(\theta) + \tilde{s}^2(\theta)}, \\ \cos \phi_c(\theta) &= \frac{\tilde{d}(\theta)}{r(\theta)}, \quad \sin \phi_c(\theta) = -\frac{\tilde{s}(\theta)}{r(\theta)}, \end{aligned} \quad (6.9)$$

with

$$\begin{aligned} \tilde{d}(\theta) &= 2(D(\theta)|\psi_d| + S(\theta)|\psi_s| \cos \phi_{sd})|\psi_{Pb}|, \\ \tilde{s}(\theta) &= 2S(\theta)|\psi_s||\psi_{Pb}| \sin \phi_{sd}. \end{aligned} \quad (6.10)$$

Combining Eqs. (4.9) and (6.8) gives, for an untwinned single crystal, a Josephson current in the standard form of Eq. (1.1), with the characteristic phase of the junction $\phi_c(\theta)$ given by Eq. (6.9). For the time-reversal invariant $d + s$ state, $\phi_{sd} = 0$ and the characteristic phase of the junction will be either zero or π . Results appropriate to a tetragonal superconductor in a $d_{x^2-y^2}$ -wave state are obtained from the above formula by taking $S(\theta) = 0$ and $D(\theta + \frac{1}{2}\pi) = -D(\theta)$. This gives $\phi_c(\theta + \frac{1}{2}\pi) = \phi_c(\theta) + \pi$ and an offset flux in a corner SQUID experiment of $\Phi_{\text{offset}} = \frac{1}{2}\Phi_0$. For an orthorhombic crystal such as YBCO, there is however no symmetry-required relationship between $D(\theta + \frac{1}{2}\pi)$ and $D(\theta)$, and also $S(\theta)$ is nonzero in general. Nevertheless, provided the quantities $\tilde{d}(\theta + \frac{1}{2}\pi)$ and $\tilde{d}(\theta)$ have appropriate signs, the $d + s$ state ($\phi_{sd} = 0$) will yield the value $\Phi_{\text{offset}} = \frac{1}{2}\Phi_0$ in a corner SQUID experiment. Under these conditions, the superconducting state of orthorhombic YBCO may be said to exhibit predominantly $d_{x^2-y^2}$ -wave behavior. For the time-reversal symmetry-breaking $d + e^{i\phi}s$ state, $\phi_{sd} \neq 0$, and $\phi_c(\theta)$ and $\phi_c(\theta + \frac{1}{2}\pi)$ differ from zero and π , and Φ_{offset} differs from zero or $\frac{1}{2}\Phi_0$. If $|\psi_s| \ll |\psi_d|$ and/or $\phi_{sd} \ll \frac{1}{2}\pi$, this difference may be small and difficult to detect experimentally.

To describe basal-plane tunneling between a twinned sample of YBCO and lead, the interface free energy f_I is averaged over the two types of twins. The interface free energy $f_{I,\bar{2}}(\theta)$ of twin 2, cf. Eq. (6.6), with the superconducting state $\Psi_{\bar{2}}$, given by Eq. (2.11), can be expressed in the form of Eq. (6.8):

$$f_{I,\bar{2}} = -r(\theta + \pi/2) \cos(\pi + \phi + \phi_c(\theta + \pi/2)), \quad (6.11)$$

which yields for the average $\bar{f}_I(\theta) = (f_{I,1} + f_{I,\bar{2}})/2$

$$\bar{f}_I(\theta) = -\bar{r}_2(\theta) \cos(\phi - \phi_{\bar{2}}(\theta)), \quad (6.12)$$

where

$$\begin{aligned} \bar{r}_2(\theta) &= \sqrt{\tilde{d}_d^2(\theta) + \tilde{s}_s^2(\theta)}, \\ \cos \phi_{\bar{2}}(\theta) &= \frac{\tilde{d}_d(\theta)}{\bar{r}_2(\theta)}, \quad \sin \phi_{\bar{2}}(\theta) = -\frac{\tilde{s}_s(\theta)}{\bar{r}_2(\theta)}. \end{aligned} \quad (6.13)$$

Here the θ -dependent functions have been separated into even (s -wave like) and odd (d -wave like) terms under $\theta \rightarrow \theta + \pi/2$ by

$$\begin{aligned}
x(\theta) &= \frac{1}{2} [x(\theta) + x(\theta + \pi/2)] + \frac{1}{2} [x(\theta) - x(\theta + \pi/2)] \\
&\equiv x_s(\theta) + x_d(\theta),
\end{aligned} \tag{6.14}$$

where $x(\theta)$ stands for $\tilde{d}(\theta)$ or $\tilde{s}(\theta)$. Also, since the two twins have been taken to be in states Ψ_1 and Ψ_2 , the twin boundaries are what have been called above $\mathcal{T}\sigma$ twin boundaries. The above results (in particular Eq. (6.13)) show that for a twinned sample with $\mathcal{T}\sigma$ twin boundaries the characteristic phases of the Josephson junctions satisfy $\phi_2(\theta + \frac{1}{2}\pi) = -\phi_2(\theta) + \pi$. This yields an offset flux in a corner SQUID experiment of

$$\Phi_{\text{offset}} = \frac{\phi_2(\theta)}{\pi} \Phi_0 + \frac{1}{2} \Phi_0. \tag{6.15}$$

The macroscopic invariance group of a crystal with $\mathcal{T}\sigma$ twin boundaries, which is the same as that of a tetragonal crystal in a $d + is$ state, does not constrain the offset flux to be 0 or $\frac{1}{2}\pi$, in agreement with Eq. (6.15). A value of $\Phi_{\text{offset}} = (0.98 \pm 0.05)\Phi_0/2$ is obtained in experiments³ on YBCO. Thus, either the YBCO twins are not in states of broken time-reversal symmetry (i.e. $\phi_{sd} = 0$), or the time-reversal symmetry-breaking is small (more explicitly, $\tilde{s}_s \ll \tilde{d}_d$), or the twin boundaries are not of the type $\mathcal{T}\sigma$, or time-reversal domains (see below) play a role.

Alternatively, if the twin boundaries are of the odd reflection symmetry type, then twin 2 is in state Ψ_2 given by Eq. (2.10), and Eq. (6.13) holds but with $\tilde{s}_s(\theta)$ replaced by $\tilde{s}_d(\theta)$. This yields $\phi_2(\theta + \frac{1}{2}\pi) = \phi_2(\theta) + \pi$ for the critical phase of the Josephson junction, and $\Phi_{\text{offset}} = \frac{1}{2}\Phi_0$. Thus, for odd-symmetry twin boundaries, the same value of the offset flux is obtained for the time-reversal invariant $d + s$ state and for the time-reversal symmetry-breaking $d + e^{i\phi}s$ state. It is therefore not possible to determine whether or not YBCO is in a time-reversal symmetry-breaking state from the experimental results of Ref. 3 alone. These results can again be qualitatively understood in terms of the macroscopic invariance group of the twinned crystal, which for this case is the same, except for the absence of time reversal, as that of the $d_{x^2-y^2}$ superconducting state of a twinned crystal.

There remains the question of the existence of domains which are time-reversal images of one another. Given a time-reversal symmetry-breaking state of a twinned crystal, a different state of the crystal having the same free energy can be obtained by applying the time-reversal operation. The various equations describing this state can be obtained from those presented in this and the preceding Section by reversing the sign of ϕ_{sd} . It may be that these two different states exist in different regions of the same crystal, in which case the crystal is said to contain time-reversal domains. If these time-reversal domains are sufficiently small, so that it is necessary to average over them to obtain the Josephson currents through a junction, then the behavior characteristic of broken time-reversal symmetry will disappear. This makes sense since a crystal containing a random distribution of domains which are time-reversed images of each other is macroscopically invariant under time reversal. Thus, the c -axis Josephson current found in the case of $\mathcal{T}\sigma$ twin boundaries, cf. Eq. (5.8), vanishes when averaged over time-reversal domains (since ϕ_1 changes sign in going from a given domain to its time reversed image). Similarly, an average of the basal plane Josephson currents over time-reversal domains will yield junctions whose characteristic phases can only be either 0 or π .

VII. PROPOSED GEOMETRY FOR RING EXPERIMENT

Recently, a number of order parameter symmetry determinations have been carried out using tricrystal ring magnetometry^{7,8,9,10,20}. These experiments have measured the flux through a superconducting ring containing three Josephson junctions. The observation of a spontaneous magnetization of the ring corresponding to a magnetic flux through the ring of $\frac{1}{2}\Phi_0$, where Φ_0 is the elementary flux quantum, shows that one or perhaps three of the Josephson junctions have a characteristic phase of π . These experiments are impressive in eliminating the possible influence of spurious trapped flux, and in the precision with which the spontaneous flux is measured.

The identification of a Josephson junction having a characteristic phase of π was taken to be evidence for $d_{x^2-y^2}$ symmetry. This conclusion was based on a simplified model³⁷ of the angular dependence (on the angles between the Josephson junction and the crystallographic axes) of the Josephson current expected for $d_{x^2-y^2}$ superconductivity. The use of this simplified model is somewhat analogous to the description of an electronic band structure in a tight binding model restricted to nearest neighbor interactions only. Such a model is a good first guess, but its reliability is in fact unknown. A complete parametrization of the Josephson current is given in Appendix B, where it is seen that, even to “lowest order”, the parametrization of Ref. 37 does not retain all terms. It should also be stated that, in its study of YBCO, Ref. 8 examined three different geometries all of which provided results consistent with the simplified d -wave model, and inconsistent with other possibilities which they raised. For the geometries studied in Refs. 7, 8, 9, 10, 20, however, it would be extremely difficult to reach definite conclusions concerning the symmetry if a general model (similar to that developed in Appendix B) for the Josephson current were adopted. For this reason,

we propose below a different ring geometry which can distinguish between $d_{x^2-y^2}$ and s wave superconductivity on the basis of symmetry arguments only, and without reference to simplified models of the Josephson junction.

It should also be noted that, as was the case with the corner SQUID experiments discussed above, the symmetry which is determined in these ring experiments on twinned crystals of YBCO is the *macroscopic* symmetry. Thus, a determination that the macroscopic symmetry is $d_{x^2-y^2}$ would lead directly to the conclusion that the twin boundaries have odd reflection symmetry, while further arguments are needed to determine the *microscopic* symmetry of the order parameter, cf. Section III C. Experiments on tetragonal crystals, such as those of Ref. 20 on $\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$ are thus of special importance in that the complicating effect of twinning is eliminated.

In view of the above, it would be useful to have an experimental test for s -wave versus $d_{x^2-y^2}$ -wave superconductivity in tetragonal crystals which is dependent on symmetry arguments only. The experimental geometry of Fig. 5 provides such a test. The superconductors labeled $S1$ and $S2$ are the same (e.g. $\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$), but $S2$ has its a and b axes oriented along the x and y axes of the figure, whereas the a and b axes of $S1$ are rotated from x and y through an angle of $\pi/4$ about the c axis.

Considerations of gauge invariance and time-reversal symmetry show that the sum of the free energies of the Josephson junctions A and B (to terms quadratic in the order parameter) has the form

$$F = c_A(\Psi_{1A}\Psi_{2A}^* + \Psi_{1A}^*\Psi_{2A}) + c_B(\Psi_{1B}\Psi_{2B}^* + \Psi_{1B}^*\Psi_{2B}), \quad (7.1)$$

where Ψ_{1A} is the order parameter of superconductor $S1$ at Josephson junction A , etc., and where c_A and c_B are real. If the superconductivity is s -wave, the invariance of the free energy with respect to rotations of the ring by π about the x axis yields the result $c_A = c_B$; in this case both junctions are either π junctions or normal junctions, and the ring will not exhibit a spontaneous magnetic moment in its ground state. On the other hand, if the superconductivity is d -wave, the same symmetry argument requires that $c_A = -c_B$; hence in this case one junction is normal while the other is a π junction and the ring will exhibit a spontaneous moment in its ground state. The flux associated with this moment is $\frac{1}{2}\Phi_0$ in the limit $(LI_c/c) \gg \Phi_0/(2\pi)$, where I_c is the critical current and L is the self-inductance of the ring^{17,20,33}.

Note that in the geometry of Fig. 5 both $\alpha = 0$ and $\alpha = \pi/4$ yield $c_A = c_B = 0$ for $d_{x^2-y^2}$ -wave superconductivity so that these angles would not be appropriate choices for the experiment. Hence, values of α such as $\alpha \sim \pi/8$ should allow a clear distinction to be made, on symmetry grounds only, between s -wave and $d_{x^2-y^2}$ -wave superconductivity.

In addition, it has been pointed out to us³⁸ that the model of Ref. 8 for the strong disordered limit of the Josephson junction yields $c_A = c_B = 0$ for all α ; thus, it would be prudent to have the least possible disorder in the junction.

We now give a detailed analysis which includes three possible types of superconductivity for a tetragonal system, cf. Section II, namely $d_{x^2-y^2}$, s and $d+is$. Assume that in general the superconductivity can be described in terms of two complex order parameters, ψ_d and ψ_s , of $d_{x^2-y^2}$ and s symmetry, respectively. Then the free energy of junction A has the form

$$F_A = G_d\psi_{d1A}\psi_{d2A}^* + G_s\psi_{s1A}\psi_{s2A}^* + G_{sd}\psi_{s1A}\psi_{d2A}^* + G_{ds}\psi_{d1A}\psi_{s2A}^* + \text{c.c.}, \quad (7.2)$$

where ψ_{d1A} is the order parameter of $d_{x^2-y^2}$ symmetry in superconductor $S1$ at junction A , etc., and where all coefficients G are real. The free energy for junction B is the same, except that $G_d \rightarrow -G_d$, $G_{ds} \rightarrow -G_{ds}$, and the subscript A is replaced by subscript B everywhere. For the $d+is$ state, the order parameter at junction A takes the form

$$\begin{aligned} \psi_{d1A} &= |\psi_{d1A}|e^{i\phi_{d1A}}, \psi_{d2A} = |\psi_{d2A}|e^{i\phi_{d2A}}, \\ \psi_{s1A} &= |\psi_{s1A}|e^{i(\phi_{d1A} + \frac{1}{2}\pi)}, \psi_{s2A} = \lambda|\psi_{s2A}|e^{i(\phi_{d2A} + \frac{1}{2}\pi)}, \end{aligned} \quad (7.3)$$

where $\lambda = \pm 1$ accounts for the freedom to choose either of the two $d \pm is$ states in superconductor $S2$. The order parameter at junction B is similarly expressed. Further analysis shows that the sum of the free energies of junctions A and B can be written as

$$F = -\frac{\Phi_0}{2\pi c}I_A \cos(\phi_A - \phi_{cA}) - \frac{\Phi_0}{2\pi c}I_B \cos(\phi_B - \phi_{cB}), \quad (7.4)$$

where I_A, I_B are positive. For pure $d_{x^2-y^2}$ -wave superconductivity, $\phi_{cA} - \phi_{cB} = \pm\pi$. For pure s -wave superconductivity, $\phi_{cA} - \phi_{cB} = 0$. For $d+is$ superconductivity, ϕ_{cA} and ϕ_{cB} are (as a result of nonzero G_{sd} and G_{ds}) different from zero (mod π). The quantities $\phi_A = \phi_{d1A} - \phi_{d2A}$ and $\phi_B = \phi_{d1B} - \phi_{d2B}$, when generalized to the usual gauge invariant form, yield

$$\frac{1}{2}(\phi_A - \phi_B) = \frac{\pi\Phi}{\Phi_0}, \quad (7.5)$$

where Φ is the flux through the ring. The quantity $\frac{1}{2}(\phi_A + \phi_B)$ is at present undetermined; minimizing F with respect to it yields

$$F = -\frac{\Phi_0}{2\pi c}(I_A + I_B) \left\{ \epsilon^2 + (1 - \epsilon^2) \cos^2 \left[\frac{\pi}{\Phi_0}(\Phi - \Phi_{\text{offset}}) \right] \right\}^{1/2} + \frac{1}{2c^2}LI^2, \quad (7.6)$$

where

$$\Phi_{\text{offset}} = \frac{\phi_{cA} - \phi_{cB}}{2\pi} \Phi_0, \quad (7.7)$$

and

$$\epsilon = \frac{|I_A - I_B|}{I_A + I_B}. \quad (7.8)$$

Here, a quantity $LI^2/(2c^2)$, representing the magnetic field energy associated with the current I circulating in the ring, has been added to F . The flux through the ring due to this current is $\Phi_I = LI/c$, where L is the ring self-inductance, and the total flux through the ring is $\Phi = \Phi_I + \Phi_{\text{ex}}$, where Φ_{ex} is the flux due to external sources.

The ground state free energy of the ring in zero external flux, $\Phi_{\text{ex}} = 0$, is found by minimizing F with respect to Φ . For sufficiently large values of the parameter $L(I_A + I_B)2\pi/(c\Phi_0)$, the term in LI^2 can be treated in perturbation theory. Also, without loss of generality, we can assume $0 \leq (\phi_{cA} - \phi_{cB}) \leq \pi$. From these assumptions it follows that the flux through the ring in its ground state is Φ_{offset} . Thus, the flux through the ring in its ground state is zero for an s -wave superconductor, $\frac{1}{2}\Phi_0$ for a $d_{x^2-y^2}$ -wave superconductor, and $x\Phi_0$, where $0 < x < \frac{1}{2}$, for a $d + is$ -wave superconductor. It should be noted that the minimum flux of $x\Phi_0$ in the ground state for the $d + is$ state occurs for the flux in a particular direction; the minimum flux in the other direction is $(1 - x)\Phi_0$, this asymmetry being a reflection of the broken time-reversal symmetry. (It has been noted previously in Ref. 8 that the $d + is$ state would lead to a ground state flux in a ring of $x\Phi_0$ where x is neither 0 nor $\frac{1}{2}$.) If the parameter $L(I_A + I_B)2\pi/(\Phi_0 c)$ is not large, the ground state flux through the ring will be less than $\frac{1}{2}\Phi_0$ for a $d_{x^2-y^2}$ superconductor, as in the case of a single-junction ring^{20,33}, so that care will have to be taken to distinguish experimentally between this situation, and the one where the departure of the ground state flux from $\frac{1}{2}\Phi_0$ is due to broken time-reversal symmetry. These results, for the geometry of Fig. 5, depend on symmetry arguments only, and do not rely for their validity on elementary models of the angular dependence of the Josephson critical currents.

Finally, note that although the observation of a ring containing flux $\frac{1}{2}\Phi_0$ in the proposed geometry would definitively rule out s -wave symmetry for the superconductivity, and would be consistent with $d_{x^2-y^2}$ -wave superconductivity, it would not definitively establish the superconductivity as being $d_{x^2-y^2}$ -wave; d_{xy} superconductivity would also be consistent with such a result. Other arguments, analogous to those of the Appendix, could however definitively establish a $d_{x^2-y^2}$ state.

VIII. CONCLUSIONS

Since YBCO is orthorhombic, the order parameter describing its superconductivity is expected to transform according to one of the irreducible representations of the orthorhombic point group. The experimental observation^{1,2,3,4,5,6,7,8,9,10,11,12,13,14} of nonzero bilinear Josephson critical currents along at least two of the three principal orthorhombic axes in tunneling from YBCO to a conventional s -wave superconductor is sufficient to identify the symmetry of the order parameter as being of the $ux^2 + vy^2$ (A_{1g}) type (see Appendix). This provides the experimental justification for considering YBCO superconductivity as coming from a mixture of $d_{x^2-y^2}$ -wave and s -wave contributions, which are the two tetragonal symmetry types compatible with orthorhombic A_{1g} symmetry.

In a model²³ of mixed s and $d_{x^2-y^2}$ superconductivity in YBCO, there is a transition to a time-reversal invariant $d + s$ state at the normal-to-superconducting transition temperature T_c , followed by a possible transition to a time-reversal symmetry-breaking $d + e^{i\phi}s$ state at a lower temperature. Corresponding to a $d + e^{i\phi}s$ state there is the time-reversed $d + e^{-i\phi}s$ state of the same free energy density so that time-reversal domains may form in a sample. Since averaging over time-reversal domains (in the case where the size of these domains is much smaller than the size of the Josephson junction) makes the effects of broken time-reversal symmetry unobservable, the case of a single time-reversal domain is considered here.

Much of this paper studies Josephson tunneling from the $d + e^{i\phi}s$ state to a conventional s -wave superconductor, and compares it with tunneling between a $d + s$ state and an s -wave superconductor. The principal effect of broken time-reversal symmetry is that the Josephson junction acquires a characteristic phase which is neither zero nor π . The differences in the characteristic phases of two junctions can be measured in a SQUID experiment.

The symmetry characteristics of the Josephson currents in a twinned crystal can be understood in terms of the macroscopic invariance group of the superconducting state of the twinned crystal. For the case where the twins are in their time-reversal invariant $d + s$ state (studied in Ref. 18) the macroscopic invariance group is that of a tetragonal $d_{x^2-y^2}$ superconductor for odd-symmetry twin boundaries, and it is that of a tetragonal s -wave superconductor for even-symmetry twin boundaries. Since macroscopic d -wave behavior is observed in basal-plane Josephson tunneling experiments on highly twinned crystals^{1,3,4,5,6,7,8,9,10}, the twin boundaries for the $d + s$ state have to be of the odd-symmetry type¹⁸, which implies that the d -wave component of the order parameter is approximately continuous across twin boundaries.

In the case of time-reversal symmetry-breaking $d + e^{i\phi}s$ states, two types of twin boundaries are consistent with a continuous d -wave component, namely the odd-symmetry and the $\mathcal{T}\sigma$ twin boundaries. For odd-symmetry twin boundaries the macroscopic invariance group is that of a tetragonal $d_{x^2-y^2}$ superconductor, but without the time-reversal operation. Detailed calculations confirm, that the Josephson tunneling for the $d + s$ state and the $d + e^{i\phi}s$ states are indistinguishable, and that they have the predicted macroscopic d -wave characteristics when the twins are present with equal weights. For $\mathcal{T}\sigma$ twin boundaries, on the other hand, the macroscopic invariance group is that of a tetragonal $d + is$ superconductor. As expected from the macroscopic symmetry, in this case the c -axis Josephson currents cancel only partially when averaged over twins, and the Josephson junction characteristic phase difference measured in a corner SQUID experiment can differ from either zero or π .

In order to determine the symmetry of the order parameter describing superconductivity in a single twin (or equivalently, the invariance group of this superconducting state), it is essential to do experiments on untwinned single crystals. Experiments on twinned crystals determine the macroscopic invariance group of the twinned crystal, which in general will be different from that of the superconducting state of an untwinned crystal, and which is determined by the symmetry of the twin boundaries.

A final section of the paper, stimulated by recent tricrystal-ring magnetometry experiments²⁰ on tetragonal $\text{Ti}_2\text{Ba}_2\text{CuO}_{6+\delta}$, proposes and analyzes a ring geometry which allows s and $d_{x^2-y^2}$ superconductivity to be distinguished by symmetry arguments only, and which makes unnecessary any assumption that an s -wave superconductor is unlikely to produce a characteristic phase of π in a Josephson junction. The characteristics of $d + is$ superconductivity for this geometry are also identified.

ACKNOWLEDGMENTS

This work was supported by the Natural Sciences and Engineering Council of Canada. Useful discussions with R. C. Dynes, A. E. Jacobs, and C. C. Tsuei are also acknowledged.

APPENDIX A: ORDER PARAMETER SYMMETRY IN YBCO

On the basis of experimental evidence, this section justifies the assumption that the superconducting order parameter in YBCO is a mixture of order parameters having $d_{x^2-y^2}$ and s wave symmetry. It is known^{17,37,39,40,41,42} that the dc Josephson effect can be used to determine the symmetry of the superconducting order parameter. Since YBCO is orthorhombic, the logical first question is: Which of the irreducible representations of the orthorhombic point group does the order parameter belong to? The argument is straightforward and elementary but we have been unable to find it in detail elsewhere and therefore present it here.

Consider a Josephson junction between YBCO and lead, the two superconductors being characterized by order parameters ψ_Γ and ψ_{Pb} , respectively. Here, ψ_Γ transforms according to the irreducible representation Γ of the orthorhombic point group D_{2h} of YBCO and ψ_{Pb} is assumed invariant under all transformations. Symmetry considerations show that the lowest order (in ψ_Γ and ψ_{Pb}) contribution to the Josephson current across a junction will have the form

$$J_{\mathbf{n}}^\Gamma = iC_{\mathbf{n}}^\Gamma [\psi_\Gamma \psi_{\text{Pb}}^* - \psi_\Gamma^* \psi_{\text{Pb}}], \quad (\text{A1})$$

where $\mathbf{n}$ is the normal to the plane of the junction, and where $J_{\mathbf{n}}$ is the current in direction $\mathbf{n}$.

Table I shows which of the coefficients $C_{\mathbf{n}}^{\Gamma}$ are zero (by symmetry) and which can be nonzero, for each irreducible representation and for currents along each of the principal orthorhombic axes. Since only the $ux^2 + vy^2$ representation in Table I can have nonzero Josephson currents along all three principal axes, and since nonzero Josephson currents have apparently been observed experimentally along all three principal axes in untwinned single crystal experiments^{1,2,13}, the $ux^2 + vy^2$ representation, also commonly called the A_{1g} representation, would appear to be the only acceptable one.

When the bilinear contribution to the Josephson current is zero, however, the biquadratic contribution

$$J_{\mathbf{n}}^{(4)\Gamma} = iC_{\mathbf{n}}^{(4)\Gamma} \left[\psi_{\Gamma}^2 \psi_{\text{Pb}}^{2*} - \psi_{\Gamma}^{2*} \psi_{\text{Pb}}^2 \right] \quad (\text{A2})$$

may be nonzero. Here, $C_{\mathbf{n}}^{(4)\Gamma}$ is never required to be zero by symmetry. Close to the critical temperature $T_{\text{c,Pb}} \simeq 7.2\text{K}$ for lead, the biquadratic critical current should have a temperature dependence $J^{(4)\Gamma} \propto |\psi_{\text{Pb}}|^2 \propto (T_{\text{c,Pb}} - T)$, i.e. linear in $T_{\text{c,Pb}} - T$, whereas the bilinear critical current should have the square root temperature dependence $J^{\Gamma} \propto |\psi_{\text{Pb}}| \propto (T_{\text{c,Pb}} - T)^{1/2}$. Measurements of the temperature dependence of the c -axis critical current by Dynes and coworkers^{11,12,13} clearly show the square root temperature dependence of the bilinear current.

We are not aware of measurements which establish experimentally that the Josephson currents along the a and b axes are of the bilinear, as opposed to the biquadratic, type. An experiment which does this would be of interest.

Note that the biquadratic Josephson current is proportional to $\sin[2(\phi_{\Gamma} - \phi_{\text{Pb}})]$, where $\phi_{\Gamma} - \phi_{\text{Pb}}$ is the phase difference between the two superconductors, whereas the bilinear Josephson current is proportional to $\sin(\phi_{\Gamma} - \phi_{\text{Pb}})$. This difference means that a SQUID made from Josephson junctions for which the current is biquadratic would have a maximum current periodic in the flux with period $\Phi_0/2$ rather than Φ_0 as is usually the case (Φ_0 is the elementary flux quantum). Also, the base frequency for the ac Josephson effect would be $\omega_J = 4eV/\hbar$ rather than $2eV/\hbar$. For a microscopic calculation of the effect for tetragonal systems see Refs. 43, 44.

If the bilinear character of the Josephson currents along the a and b axes of YBCO can be confirmed experimentally, then the $ux^2 + vy^2$ character of the superconducting order parameter in YBCO will be established. Of all the different irreducible representations of the tetragonal point group, only the $d_{x^2-y^2}$ and s representations are invariant under the operations of the orthorhombic point group. (A different way of seeing this is to note that $ux^2 + vy^2 = d(x^2 - y^2) + s(x^2 + y^2)$, where the coefficients d and s can be found in terms of a and b .) Thus the $ux^2 + vy^2$ order parameter for YBCO can be considered to be a combination of $d_{x^2-y^2}$ and s (e.g. $x^2 + y^2$) basis vectors (and no other type of tetragonal symmetry). These arguments assume a second order transition to the superconducting state.

APPENDIX B: ANGULAR DEPENDENCE OF JOSEPHSON CURRENTS

Figure 6 shows a Josephson junction between two identical tetragonal superconductors. The free energy per unit area of the Josephson junction in zero magnetic field has the form

$$F = C(\theta, \theta') \cos(\phi - \phi'), \quad (\text{B1})$$

where the superconducting order parameters on the two sides of the junction are $\psi = |\psi|e^{i\phi}$ and $\psi' = |\psi'|e^{i\phi'}$, and θ and θ' define the orientations of the crystallographic axes, as shown in the figure.

First assume that the superconductivity is either s -wave or $d_{x^2-y^2}$ -wave. (Our definition of s -wave is that the order parameter is invariant with respect to all operations of the tetragonal point group; this is sometimes called generalized s -wave superconductivity. Similarly, our definition of $d_{x^2-y^2}$ -wave superconductivity encompasses generalized $d_{x^2-y^2}$ -wave superconductivity.) Then symmetry considerations show that the coefficients must satisfy the following conditions:

$$C(\theta, \theta') = C(\theta', \theta) = C(-\theta, -\theta') = C(\theta + \pi, \theta'). \quad (\text{B2})$$

Furthermore, for s -wave superconductivity

$$C_s(\theta + \pi/2, \theta') = C_s(\theta, \theta'), \quad (\text{B3})$$

whereas for $d_{x^2-y^2}$ superconductivity

$$C_d(\theta + \pi/2, \theta') = -C_d(\theta, \theta'). \quad (\text{B4})$$

The use of these constraints yields the following Fourier series representations for the coefficients:

$$C_s(\theta, \theta') = \sum_{n, n'} [C_{4n, 4n'} \cos(4n\theta) \cos(4n'\theta') + S_{4n, 4n'} \sin(4n\theta) \sin(4n'\theta')] \quad (\text{B5})$$

for s -wave superconductivity, and

$$C_d(\theta, \theta') = \sum_{n, n'} \{C_{4n+2, 4n'+2} \cos[(4n+2)\theta] \cos[(4n'+2)\theta'] + S_{4n+2, 4n'+2} \sin[(4n+2)\theta] \sin[(4n'+2)\theta']\} \quad (\text{B6})$$

for $d_{x^2-y^2}$ -wave superconductivity. The sums over n and n' are over all positive integers and zero; also, the coefficients have the property $C_{i,j} = C_{j,i}$ and $S_{i,j} = S_{j,i}$, where i, j are $4n$, $4n'$ or $4n+2$, $4n'+2$.

The first few terms of these expressions are

$$C_s(\theta, \theta') = C_{0,0} + C_{4,0} [\cos(4\theta) + \cos(4\theta')] + \dots, \quad (\text{B7})$$

and

$$C_d(\theta, \theta') = C_{2,2} \cos(2\theta) \cos(2\theta') + S_{2,2} \sin(2\theta) \sin(2\theta') + \dots \quad (\text{B8})$$

Note that the expression proposed by Sigrist and Rice³⁷ for the $d_{x^2-y^2}$ case, namely $C_{\text{SR}}(\theta, \theta') = C_{2,2} \cos(2\theta) \cos(2\theta')$, has the property $C_{\text{SR}}(\theta, \pi/4) = 0$, which is not required by symmetry and therefore cannot be expected to be true in general. The simplified expression given in Eq. (B8), which contains two constants $C_{2,2}$ and $S_{2,2}$, does not have this difficulty. Also, if $S_{2,2} = -C_{2,2}$, the sines and cosines in our two-term expression Eq. (B8) can be summed to give $C_d = C_{2,2} \cos[2(\theta + \theta')]$, which is the expression proposed by Tsuei *et al.*^{8,20} to model what they call the maximum disorder limit of a grain boundary Josephson junction.

- ¹ D. A. Wollman, D. J. Van Harlingen, W. C. Lee, D. M. Ginsberg, and A. J. Leggett, Phys. Rev. Lett. **71**, 2134 (1993).
- ² D. A. Brawner and H. R. Ott, Phys. Rev. B **50**, 6530 (1994), D. Brawner, private communication.
- ³ A. Mathai, Y. Gim, R. C. Black, A. Amar, and F. C. Wellstood, Phys. Rev. Lett. **74**, 4523 (1995).
- ⁴ D. A. Wollman, D. J. Van Harlingen, J. Giapintzakis, and D. M. Ginsberg, Phys. Rev. Lett. **74**, 797 (1995).
- ⁵ D. J. van Harlingen, Rev. Mod. Phys. **67**, 515 (1995).
- ⁶ I. Iguchi and Z. Wen, Phys. Rev. B **49**, 12388 (1994).
- ⁷ C. C. Tsuei, J. R. Kirtley, C. C. Chi, L. S. Yu-Jahnes, A. Gupta, T. Shaw, J. Z. Sun, and M. B. Ketchen, Phys. Rev. Lett. **73**, 593 (1994).
- ⁸ C. C. Tsuei, J. R. Kirtley, M. Rupp, L. S. Yu-Jahnes, and C. C. Chi, J. Phys. Chem. Solids **56**, 1787 (1995).
- ⁹ C. C. Tsuei, D. M. Newns, J. R. Kirtley, and P. C. Pattnaik, in *High T_c Superconductivity and the C_{60} Family*, edited by S. Feng and H.-C. Ren (Gordon and Breach, 1995), pp. 293–324.
- ¹⁰ J. R. Kirtley, C. C. Tsuei, M. Rupp, J. Z. Sun, L. S. Yu-Jahnes, A. Gupta, M. B. Ketchen, K. A. Moler, and M. Bhushan, preprint, 1995.
- ¹¹ A. G. Sun, D. A. Gajewski, M. B. Maple, and R. C. Dynes, Phys. Rev. Lett. **72**, 2267 (1994).
- ¹² A. S. Katz, A. G. Sun, and R. C. Dynes, Appl. Phys. Lett. **66**, 105 (1995).
- ¹³ R. C. Dynes, Physica C **235–240**, 225 (1994).
- ¹⁴ D. A. Wollman, D. J. Van Harlingen, D. M. Ginsberg, and J. Giapintzakis, preprint presented at Stanford Conf. on Novel Spectroscopies, 1995.
- ¹⁵ P. Chaudhari and S.-Y. Lin, Phys. Rev. Lett. **72**, 1084 (1994).
- ¹⁶ E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994).
- ¹⁷ M. Sigrist and T. M. Rice, Rev. Mod. Phys. **67**, 503 (1995).
- ¹⁸ M. B. Walker, Phys. Rev. B, in press, eprint cond-mat/9510013, 1995.
- ¹⁹ A. F. Andreev, JETP Lett. **46**, 584 (1987).
- ²⁰ C. C. Tsuei, J. R. Kirtley, M. Rupp, J. Z. Sun, A. Gupta, and M. B. Ketchen, Science **271**, 329 (1996).
- ²¹ A. J. Millis, Phys. Rev. B **49**, 15408 (1994).
- ²² J. R. Kirtley, P. Chaudhari, M. B. Ketchen, N. Khare, S. Y. Lin, and T. Shaw, Phys. Rev. B **51**, 12057 (1995).
- ²³ Q. P. Li, E. C. Koltenbah, and R. Joynt, Phys. Rev. B **48**, 437 (1993).
- ²⁴ Y. Itoh, Y. Yoshimura, T. Ohomura, H. Yasuoka, Y. Ueda, and K. Kosuge, J. Phys. Soc. Jpn. **63**, 1455 (1994).

- ²⁵ J.-H. Xu, Y. Ren, and C. S. Ting, eprint: cond-mat/9511130, 1995.
- ²⁶ S. Spielman, K. Fesler, C. B. Eom, T. H. Geballe, M. M. Fejer, and A. Kapitulnik, Phys. Rev. Lett. **65**, 123 (1990).
- ²⁷ T. W. Lawrence, A. Szöke, and R. B. Laughlin, Phys. Rev. Lett. **69**, 1439 (1992).
- ²⁸ R. F. Kiefl, J. H. Brewer, I. Affleck, J. F. Carolan, P. Dosanjih, W. N. Hardy, T. Hsu, R. Kadono, J. R. Kempton, S. R. Kreitzman, Q. Li, A. H. O'Reilly, T. M. Riseman, P. Schleger, P. C. E. Stamp, H. Zhou, P. G. Le, G. M. Luke, B. Sternlieb, Y. J. Uemura, H. R. Hart, and K. W. Lay, Phys. Rev. Lett. **64**, 2082 (1990).
- ²⁹ M. Sigrist, D. B. Bailey, and R. B. Laughlin, Phys. Rev. Lett. **74**, 3249 (1995).
- ³⁰ M. Sigrist, K. Kuboki, P. A. Lee, A. J. Millis, and T. M. Rice, Phys. Rev. B **53**, 2835 (1996).
- ³¹ Y. Zhu, M. Suenaga, and J. Taftø, Philosophical Magazine A **67**, 1057 (1993).
- ³² G. E. Volovik and L. P. Gor'kov, Sov. Phys. JETP **61**, 843 (1985).
- ³³ L. N. Bulaevskii, V. V. Kuzii, and A. A. Sobyenin, JETP Lett. **25**, 290 (1977).
- ³⁴ D. N. Basov, R. Liang, D. A. Bonn, W. N. Hardy, B. Dabrowski, M. Quijada, D. B. Tanner, J. P. Rice, D. M. Ginsberg, and T. Timusk, Phys. Rev. Lett. **74**, 598 (1995).
- ³⁵ A. G. Sun, S. H. Han, A. S. Katz, D. A. Gajewski, M. B. Maple, and R. C. Dynes, Phys. Rev. B **52**, 15731 (1995).
- ³⁶ W. A. Atkinson and J. P. Carbotte, Phys. Rev. B **52**, 10601 (1995).
- ³⁷ M. Sigrist and T. M. Rice, J. Phys. Soc. Jpn. **61**, 4283 (1992).
- ³⁸ C. C. Tsuei, private communication.
- ³⁹ V. B. Geshkenbein and A. I. Larkin, JETP Lett. **43**, 395 (1986).
- ⁴⁰ V. B. Geshkenbein, A. I. Larkin, and A. Barone, Phys. Rev. B **36**, 235 (1987).
- ⁴¹ L. P. Gor'kov, Sov. Sci. Rev. A Phys. **9**, 1 (1987).
- ⁴² J. F. Annett, Adv. Phys. **39**, 83 (1990).
- ⁴³ Y. Tanaka, Phys. Rev. Lett. **72**, 3871 (1994).
- ⁴⁴ W. Zhang, Phys. Rev. B **52**, 3772 (1995).

TABLE I. Josephson Currents for Orthorhombic Symmetry. The value of the critical Josephson current is given for each irreducible representation Γ and for each of the orthorhombic principal axes, a , b , and c .

Γ	a	b	c
$ux^2 + vy^2$	C_a^1	C_b^1	C_c^1
yz	0	0	0
zx	0	0	0
xy	0	0	0
xyz	0	0	0
x	C_a^6	0	0
y	0	C_b^7	0
z	0	0	C_c^8

FIG. 1. Unit cells in neighboring twins under the assumption that the twin boundary (dashed line) is a plane of reflection symmetry of the crystal.


FIG. 2. Schematic representation of $d_{x^2-y^2}$ - and s -wave order-parameter components near odd and even reflection-symmetry twin-boundaries.



FIG. 3. (a) The structure of the interface between a region containing (110) twin boundaries and a region containing ($\bar{1}10$) twin boundaries, as described in Ref. 31. The rectangles indicate the relative orientations of the orthorhombic a and b axes in the different twins, and the plus and minus signs indicate the relative sign of the superconducting order parameter in the different twins. The c axis is normal to the plane of the figure. Part (b) of the figure is obtained from part (a) by a rotation of $\pi/2$ about the c -axis.

FIG. 4. Corner SQUID geometry for basal-plane tunneling between YBCO and lead. A and B indicate basal-plane Josephson junctions on two perpendicular faces of the YBCO crystal, while I_A and I_B denote the corresponding currents.

FIG. 5. Proposed geometry for ring experiment to test for s -wave versus $d_{x^2-y^2}$ -wave superconductivity in tetragonal systems. $S1$ and $S2$ are tetragonal superconductors with their c -axis normal to the plane of the figure and their basal-plane unit cells rotated by an angle $\pi/4$ with respect to each other. A and B are Josephson junctions; the angle α is chosen so that $\alpha \neq 0$, $\alpha \neq \pi/4$.

FIG. 6. Josephson junction between two identical tetragonal superconductors.






(110) twin
boundaries

(-110) twin
boundaries

(a)



(b)

